





EAST Search History

Ref #	Hits	Search Query	DBs	Default Operator	Plurals	Time Stamp
L1	207	piribedil	US-PGPUB; USPAT; USOCR; EPO; JPO; DERWENT	OR	ON	2006/08/21 16:19
L2	1722	(544/295).CCLS.	US-PGPUB; USPAT; USOCR; EPO; JPO; DERWENT	OR	OFF	2006/08/21 16:19
L3	1926	1 or 2	US-PGPUB; USPAT; USOCR; EPO; JPO; DERWENT	OR	ON	2006/08/21 16:19

NPL

		Results
2.	TITLE-ABSTR-KEY(piribedil) and TITLE-ABSTR-KEY(purification or purity or pure) [All Sources(- All Sciences -)]	7
1.	TITLE-ABSTR-KEY(piribedil) [All Sources(- All Sciences -)]	595

Copyright © 2006 Elsevier B.V. All rights reserved.
ScienceDirect® is a registered trademark of Elsevier B.V.



chain nodes :

19 24 25 26 27 28 29

ring nodes :

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 20 21 22

chain bonds :

2-26 3-25 4-24 6-7 10-19 13-29 14-28 15-19 16-27

ring bonds :

1-2 1-6 2-3 3-4 4-5 5-6 7-8 7-12 8-9 9-10 10-11 11-12 13-14 13-18 14-15 15-16 16-17
17-18 17-20 18-22 20-21 21-22

exact/norm bonds :

6-7

exact bonds :

2-26 3-25 4-24 7-8 7-12 8-9 9-10 10-11 10-19 11-12 13-29 14-28 15-19 16-27 17-20 18-22
20-21 21-22

normalized bonds :

1-2 1-6 2-3 3-4 4-5 5-6 13-14 13-18 14-15 15-16 16-17 17-18

isolated ring systems :


containing 1 : 7 : 13 :

Match level :

1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:Atom 8:Atom 9:Atom 10:Atom 11:Atom 12:Atom
13:Atom 14:Atom 15:Atom 16:Atom 17:Atom 18:Atom 19:CLASS20:Atom 21:Atom 22:Atom 24:CLASS
25:CLASS26:CLASS27:CLASS28:CLASS29:CLASS

=>

Uploading C:\Program Files\Stnexp\Queries\10810799.str



chain nodes :

19 24 25 26 27 28 29

ring nodes :

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 20 21 22

chain bonds :

2-26 3-25 4-24 6-7 10-19 13-29 14-28 15-19 16-27

ring bonds :

1-2 1-6 2-3 3-4 4-5 5-6 7-8 7-12 8-9 9-10 10-11 11-12 13-14 13-18
14-15 15-16 16-17 17-18 17-20 18-22 20-21 21-22

exact/norm bonds :

6-7

exact bonds :

2-26 3-25 4-24 7-8 7-12 8-9 9-10 10-11 10-19 11-12 13-29 14-28 15-19
16-27 17-20 18-22 20-21 21-22

normalized bonds :

1-2 1-6 2-3 3-4 4-5 5-6 13-14 13-18 14-15 15-16 16-17 17-18

isolated ring systems :

containing 1 : 7 : 13 :

Match level :

1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:Atom 8:Atom 9:Atom 10:Atom
11:Atom 12:Atom 13:Atom 14:Atom 15:Atom 16:Atom 17:Atom 18:Atom 19:CLASS
20:Atom 21:Atom 22:Atom 24:CLASS 25:CLASS 26:CLASS 27:CLASS 28:CLASS
29:CLASS


L1 STRUCTURE UPLOADED

=> d l1

L1 HAS NO ANSWERS

L1 STR

10/810,799



Structure attributes must be viewed using STN Express query preparation.

=> s l1 sss sam

SAMPLE SEARCH INITIATED 15:26:09 FILE 'REGISTRY'

SAMPLE SCREEN SEARCH COMPLETED - 12 TO ITERATE

100.0% PROCESSED 12 ITERATIONS

1 ANSWERS

SEARCH TIME: 00.00.01

FULL FILE PROJECTIONS: ONLINE **COMPLETE**

BATCH **COMPLETE**

PROJECTED ITERATIONS: 33 TO 447

PROJECTED ANSWERS: 1 TO 80

L2 1 SEA SSS SAM L1

=> =>Testing the current file.... screen

ENTER SCREEN EXPRESSION OR (END):end


=> screen 2039 OR 2040 OR 2041 OR 2047 OR 2127

L3 SCREEN CREATED

=>

Uploading C:\Program Files\Stnexp\Queries\10810799 (a).str

10/810,799



chain nodes :

19 24 25 26 27 28 29

ring nodes :

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 20 21 22

chain bonds :

2-26 3-25 4-24 6-7 10-19 13-29 14-28 15-19 16-27

ring bonds :

1-2 1-6 2-3 3-4 4-5 5-6 7-8 7-12 8-9 9-10 10-11 11-12 13-14 13-18
14-15 15-16 16-17 17-18 17-20 18-22 20-21 21-22

exact/norm bonds :

6-7

exact bonds :

2-26 3-25 4-24 7-8 7-12 8-9 9-10 10-11 10-19 11-12 13-29 14-28 15-19
16-27 17-20 18-22 20-21 21-22

normalized bonds :

1-2 1-6 2-3 3-4 4-5 5-6 13-14 13-18 14-15 15-16 16-17 17-18

isolated ring systems :

containing 1 : 7 : 13 :

Match level :

1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:Atom 8:Atom 9:Atom 10:Atom
11:Atom 12:Atom 13:Atom 14:Atom 15:Atom 16:Atom 17:Atom 18:Atom 19:CLASS
20:Atom 21:Atom 22:Atom 24:CLASS 25:CLASS 26:CLASS 27:CLASS 28:CLASS
29:CLASS

L4 STRUCTURE UPLOADED

=> que L4 NOT L3


L5 QUE L4 NOT L3

=> d 15

L5 HAS NO ANSWERS

L3 SCR 2039 OR 2040 OR 2041 OR 2047 OR 2127

L4 STR



Structure attributes must be viewed using STN Express query preparation.
 L5 QUE L4 NOT L3

=> s 15 sss sam
 SAMPLE SEARCH INITIATED 15:27:44 FILE 'REGISTRY'
 SAMPLE SCREEN SEARCH COMPLETED - 9 TO ITERATE

100.0% PROCESSED 9 ITERATIONS 0 ANSWERS
 SEARCH TIME: 00.00.01

FULL FILE PROJECTIONS: ONLINE **COMPLETE**
 BATCH **COMPLETE**
 PROJECTED ITERATIONS: 9 TO 360
 PROJECTED ANSWERS: 0 TO 0

L6 0 SEA SSS SAM L4 NOT L3

=> s 15 sss ful
 FULL SEARCH INITIATED 15:28:02 FILE 'REGISTRY'
 FULL SCREEN SEARCH COMPLETED - 344 TO ITERATE

100.0% PROCESSED 344 ITERATIONS 1 ANSWERS
 SEARCH TIME: 00.00.01

L7 1 SEA SSS FUL L4 NOT L3

=> => s 17
 L8 375 L7

=> s purification
 L9 325498 PURIFICATION

=> s 18 and 19
 L10 1 L8 AND L9

=> d bib,ab,hitstr

L10 ANSWER 1 OF 1 CAPLUS COPYRIGHT 2006 ACS on STN
 AN 2005:1050937 CAPLUS
 DN 143:326397
 TI purification of Piribedil by crystallization from water/ethanol.
 IN Jong, Shean-Jeng; Lin, Yu-Sheng
 PA Chung-Shan Institute of Science & Technology, Taiwan
 SO U.S. Pat. Appl. Publ., 3 pp.
 CODEN: USXXCO

DT Patent
 LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 2005215788	A1	20050929	US 2004-810799	20040329
PRAI	US 2004-810799		20040329		


AB A method for purifying Piribedil (I) comprises mixing I solid having a purity of ≤ 98 weight% with H₂O, boiling the mixture, adding slowly 95% EtOH to the boiling mixture to form a clear liquid, filtering the hot clear liquid, and cooling the filtrate to obtain 99.8 weight% pure white crystalline

I.

IT 3605-01-4P, Piribedil
 RL: PUR (Purification or recovery); PREP (Preparation)
 (purification of piribedil by crystallization from water/ethanol)


RN 3605-01-4 CAPLUS

CN Pyrimidine, 2-[4-(1,3-benzodioxol-5-ylmethyl)-1-piperazinyl]- (9CI) (CA INDEX NAME)




```
=> s pur?  
L11      1778040 PUR?  
  
=> s l8 and l11  
L12      20 L8 AND L11  
  
=> s l12 not l10  
L13      19 L12 NOT L10  
  
=> d l13 1-19 bib,ab,hitstr
```

L13 ANSWER 1 OF 19 CAPLUS COPYRIGHT 2006 ACS on STN
 AN 2006:232815 CAPLUS
 DN 144:370053
 TI Parallel synthesis of N-arylpiperazines using polymer-assisted reactions
 AU Duncton, Matthew A. J.; Roffey, Jonathan R. A.; Hamlyn, Richard J.; Adams, David R.
 CS Department of Chemistry, Vernalis Research Ltd, Winnersh, Wokingham, RG41 5UA, UK
 SO Tetrahedron Letters (2006), 47(15), 2549-2552
 CODEN: TELEAY; ISSN: 0040-4039
 PB Elsevier B.V.
 DT Journal
 LA English
 AB A series of N-arylpiperazines were prepared in a parallel fashion using palladium-catalyzed cross-coupling, or nucleophilic aromatic displacement chemistries, and polymer-assisted sequestration and purification techniques as key steps.
 IT 3605-01-4P
 RL: CPN (Combinatorial preparation); CMBI (Combinatorial study); PREP (Preparation)
 (parallel synthesis of N-arylpiperazines using palladium-catalyzed cross-coupling or nucleophilic aromatic displacement)
 RN 3605-01-4 CAPLUS
 CN Pyrimidine, 2-[4-(1,3-benzodioxol-5-ylmethyl)-1-piperazinyl]- (9CI) (CA INDEX NAME)



RE.CNT 10 THERE ARE 10 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 2 OF 19 CAPLUS COPYRIGHT 2006 ACS on STN
 AN 2005:962085 CAPLUS
 DN 143:254025
 TI Dopamine-agonist combination therapy for improving sleep quality
 IN Barberich, Timothy J.
 PA Sepracor, Inc., USA
 SO PCT Int. Appl., 100 pp.
 CODEN: PIXXD2

DT Patent
 LA English
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2005079851	A2	20050901	WO 2005-US3937	20050207
	WO 2005079851	A3	20060622		
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW, SM			
	RW:	BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
	US 2005267176	A1	20051201	US 2005-52719	20050207
PRAI	US 2004-545413P	P	20040218		


AB The present invention generally to pharmaceutical compns. comprising a dopamine agonist and sedative agent. In a preferred embodiment, the dopamine agonist is optically pure (S)-didesmethyisibutramine (S-I). In a preferred embodiment, the sedative agent is optically pure (S)-zopiclone or optically pure (S)-N-desmethylzopiclone. In a preferred embodiment, the dopamine agonist is optically pure (S)-I; and the sedative agent is optically pure (S)-zopiclone or optically pure (S)-N-desmethylzopiclone. The pharmaceutical compns. of the invention are useful in the treatment of restless-leg syndrome and periodic-limb-movement disorder, as well as various sleep disorders. In addition, the present invention relates to a method of treating a patient suffering from restless-leg syndrome, periodic-limb-movement disorder, a sleep abnormality, or insomnia comprising coadministering a therapeutically effective amount of a dopamine agonist and a therapeutically effective amount of a sedative agent. The preparation of (S)-I L-tartrate is given.

IT 3605-01-4, Piribedil


RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (dopamine-agonist combination therapy for improving sleep quality)

RN 3605-01-4 CAPLUS

CN Pyrimidine, 2-[4-(1,3-benzodioxol-5-ylmethyl)-1-piperazinyl]- (9CI) (CA INDEX NAME)



L13 ANSWER 3 OF 19 CAPLUS COPYRIGHT 2006 ACS on STN
 AN 2004:1068075 CAPLUS
 DN 142:168975
 TI "Lead Hopping". Validation of Topomer Similarity as a Superior Predictor of Similar Biological Activities
 AU Cramer, Richard D.; Jilek, Robert J.; Guessregen, Stefan; Clark, Stephanie J.; Wendt, Bernd; Clark, Robert D.
 CS Tripos Discovery Research, Cornwall, EX23 8LY, UK
 SO Journal of Medicinal Chemistry (2004), 47(27), 6777-6791
 CODEN: JMCMAR; ISSN: 0022-2623
 PB American Chemical Society
 DT Journal
 LA English
 AB Two extensive studies quantifying the ability of topomer shape similarity to forecast a variety of biol. similarities are described. In a prospective trial of "lead hopping", using topomer similarity for virtual screening and queries from the patent literature, biol. assays of 308 selected compds. (representing 0.03% of those available, per assay type) yielded 11 successful "lead hops" in the 13 assays attempted. The hit rate averaged over all assays was 39% ("activity" defined as inhibition $\geq 20\%$ at 10 μM), significantly greater than an unexpectedly high neg. control hit rate of 15%. The average "Tanimoto 2D fingerprint similarity" between query and "lead hop" structures (0.36) was little more than the Tanimoto similarity between random drug-like structures. Topomer shape and Tanimoto 2D fingerprint similarities were also compared retrospectively, in their tendencies to concentrate together potential and actual drugs reported to belong to the same "activity class", for twenty classes. Among the most similar 3% of structures (corresponding to " ≥ 0.85 Tanimoto" for these structures), an average of 62% of the topomer similar selection possessed a near neighbor belonging to the same activity class, roughly a one-third superiority over the "Tanimoto ≥ 0.85 " selection containing 48% actives in avoiding false positives. Conversely, the least similar 75% of structures contained 0.3% actives for topomer similarity vs. 1.0% actives for Tanimoto 2D fingerprint similarity, a 3-fold superiority for topomers in avoiding false negatives.
 IT 3605-01-4, Piribedil
 RL: PAC (Pharmacological activity); BIOL (Biological study)
 (validation of topomer similarity as a superior predictor of similar biol. activities of "Lead hopping")
 RN 3605-01-4 CAPLUS
 CN Pyrimidine, 2-[4-(1,3-benzodioxol-5-ylmethyl)-1-piperazinyl]- (9CI) (CA INDEX NAME)



RE.CNT 49 THERE ARE 49 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 4 OF 19 CAPLUS COPYRIGHT 2006 ACS on STN
 AN 2004:430288 CAPLUS
 DN 140:429017
 TI Drug condensation aerosols and kits
 IN Hale, Ron L.; Hodges, Craig C.; Lloyd, Peter M.; Lu, Amy T.; Myers, Daniel J.; Rabinowitz, Joshua D.; Wensley, Martin J.
 PA Alexza Molecular Delivery Corporation, USA
 SO U.S. Pat. Appl. Publ., 84 pp., Cont.-in-part of U.S. Ser. No. 633,877.
 CODEN: USXXCO
 DT Patent
 LA English
 FAN.CNT 32

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	---	-----	-----	-----
PI	US 2004099269	A1	20040527	US 2003-718982	20031120
	US 7090830	B2	20060815		
	US 2003051728	A1	20030320	US 2001-57198	20011026
	US 2003015197	A1	20030123	US 2002-146088	20020513
	US 2003017115	A1	20030123	US 2002-146516	20020513
	US 6737042	B2	20040518		
	US 2003035776	A1	20030220	US 2002-146515	20020513
	US 6682716	B2	20040127		
	US 2003209240	A1	20031113	US 2002-146086	20020513
	US 2003007933	A1	20030109	US 2002-150267	20020515
	US 6797259	B2	20040928		
	US 2003007934	A1	20030109	US 2002-150268	20020515
	US 6780399	B2	20040824		
	US 2003091511	A1	20030515	US 2002-150056	20020515
	US 6805853	B2	20041019		
	US 2003017117	A1	20030123	US 2002-151596	20020516
	US 6855310	B2	20050215		
	US 2003206869	A1	20031106	US 2002-151626	20020516
	US 6783753	B2	20040831		
	US 2003017116	A1	20030123	US 2002-150857	20020517
	US 6716415	B2	20040406		
	US 2003021753	A1	20030130	US 2002-150591	20020517
	US 6780400	B2	20040824		
	US 2003005924	A1	20030109	US 2002-152652	20020520
	US 6740307	B2	20040525		
	US 2003012740	A1	20030116	US 2002-153139	20020520
	US 6814954	B2	20041109		
	US 2003017118	A1	20030123	US 2002-152639	20020520
	US 6716416	B2	20040406		
	US 2003021754	A1	20030130	US 2002-152640	20020520
	US 6743415	B2	20040601		
	US 2003012737	A1	20030116	US 2002-153311	20020521
	US 6884408	B2	20050426		
	US 2003015189	A1	20030123	US 2002-153831	20020521
	US 6740308	B2	20040525		
	US 2003017119	A1	20030123	US 2002-153839	20020521
	US 6776978	B2	20040817		
	US 2003032638	A1	20030213	US 2002-153313	20020521
	US 2003005925	A1	20030109	US 2002-155621	20020522
	US 6759029	B2	20040706		
	US 2003012738	A1	20030116	US 2002-155373	20020522
	US 6737043	B2	20040518		
	US 2003017120	A1	20030123	US 2002-155703	20020522
	US 6803031	B2	20041012		

US 2003021755	A1	20030130	US 2002-155705	20020522
US 6805854	B2	20041019		
US 2003000518	A1	20030102	US 2002-155097	20020523
US 6716417	B2	20040406		
US 2003015190	A1	20030123	US 2002-154594	20020523
US 6740309	B2	20040525		
US 2003017114	A1	20030123	US 2002-154765	20020523
US 6814955	B2	20041109		
US 2003118512	A1	20030626	US 2002-280315	20021025
WO 2003045484	A2	20030605	WO 2002-US37491	20021121
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
AU 2002364508	A1	20030610	AU 2002-364508	20021121
US 2003138382	A1	20030724	US 2002-302010	20021121
US 7078016	B2	20060718		
US 2003138508	A1	20030724	US 2002-322227	20021217
US 2004126326	A1	20040701	US 2003-734902	20031212
US 7029658	B2	20060418		
US 2004127481	A1	20040701	US 2003-735198	20031212
US 7008615	B2	20060307		
US 2004126327	A1	20040701	US 2003-735199	20031212
US 7070761	B2	20060704		
US 2004127490	A1	20040701	US 2003-735495	20031212
US 7018619	B2	20060328		
US 2004126328	A1	20040701	US 2003-735496	20031212
US 7005121	B2	20060228		
US 2004126329	A1	20040701	US 2003-735497	20031212
US 7070762	B2	20060704		
US 2004156788	A1	20040812	US 2003-749535	20031230
US 2004156789	A1	20040812	US 2003-749536	20031230
US 2004156790	A1	20040812	US 2003-749783	20031230
US 7078019	B2	20060718		
US 2004156791	A1	20040812	US 2003-750303	20031230
US 7078020	B2	20060718		
US 2005075273	A1	20050407	US 2003-749539	20031230
US 7078018	B2	20060718		
US 2005089479	A1	20050428	US 2003-749537	20031230
US 7078017	B2	20060718		
US 2004184996	A1	20040923	US 2004-766279	20040127
US 7087217	B2	20060808		
US 2004191179	A1	20040930	US 2004-766566	20040127
US 7060254	B2	20060613		
US 2004191180	A1	20040930	US 2004-766574	20040127
US 7045118	B2	20060516		
US 2004191181	A1	20040930	US 2004-766634	20040127
US 7070763	B2	20060704		
US 2004191182	A1	20040930	US 2004-766647	20040127
US 7070764	B2	20060704		
US 2004228807	A1	20041118	US 2004-766149	20040127
US 7087216	B2	20060808		

US 2004184997	A1	20040923	US 2004-767115	20040128
US 7052679	B2	20060530		
US 2004184998	A1	20040923	US 2004-768205	20040129
US 7070765	B2	20060704		
US 2004184999	A1	20040923	US 2004-768220	20040129
US 7063830	B2	20060620		
US 2004185000	A1	20040923	US 2004-768293	20040129
US 7067114	B2	20060627		
US 2004185003	A1	20040923	US 2004-769157	20040129
US 7060255	B2	20060613		
US 2004185004	A1	20040923	US 2004-769197	20040129
US 7063831	B2	20060620		
US 2004202617	A1	20041014	US 2004-768281	20040129
US 2004185001	A1	20040923	US 2004-769046	20040130
US 7070766	B2	20060704		
US 2004185002	A1	20040923	US 2004-769051	20040130
US 7033575	B2	20060425		
US 2004161385	A1	20040819	US 2004-775586	20040209
US 7048909	B2	20060523		
US 2004167228	A1	20040826	US 2004-775583	20040209
US 7018620	B2	20060328		
US 2004170569	A1	20040902	US 2004-791915	20040303
US 7005122	B2	20060228		
US 2004170570	A1	20040902	US 2004-792012	20040303
US 7018621	B2	20060328		
US 2004170572	A1	20040902	US 2004-792096	20040303
US 7011819	B2	20060314		
US 2004170573	A1	20040902	US 2004-792239	20040303
US 7014840	B2	20060321		
US 2004185005	A1	20040923	US 2004-813721	20040331
US 7022312	B2	20060404		
US 2004186130	A1	20040923	US 2004-813722	20040331
US 7063832	B2	20060620		
US 2004191183	A1	20040930	US 2004-814690	20040331
US 7014841	B2	20060321		
US 2004191184	A1	20040930	US 2004-814998	20040331
US 2004185006	A1	20040923	US 2004-815527	20040401
US 6994843	B2	20060207		
US 2004185007	A1	20040923	US 2004-816407	20040401
US 7011820	B2	20060314		
US 2004185008	A1	20040923	US 2004-816567	20040401
US 7052680	B2	20060530		
US 2004191185	A1	20040930	US 2004-816492	20040401
US 7008616	B2	20060307		
US 2006153779	A1	20060713	US 2006-370628	20060307
US 2006177382	A1	20060810	US 2006-398383	20060404
PRAI US 2001-57197	A2	20011026		
US 2001-57198	A2	20011026		
US 2001-345876P	P	20011109		
US 2001-345882P	P	20011109		
US 2001-332165P	P	20011121		
US 2001-332279P	P	20011121		
US 2001-332280P	P	20011121		
US 2001-342066P	P	20011218		
US 2002-50056	B2	20020114		
US 2002-57098	A2	20020123		
US 2002-371457P	P	20020409		
US 2002-146080	A2	20020513		

US 2002-146086	A2	20020513
US 2002-146088	A2	20020513
US 2002-146515	A2	20020513
US 2002-146516	A2	20020513
US 2002-150056	A2	20020515
US 2002-150267	A2	20020515
US 2002-150268	A2	20020515
US 2002-151596	A2	20020516
US 2002-151626	A2	20020516
US 2002-150591	A2	20020517
US 2002-150857	A2	20020517
US 2002-152639	A2	20020520
US 2002-152640	A2	20020520
US 2002-152652	A2	20020520
US 2002-153139	A2	20020520
US 2002-153311	A2	20020521
US 2002-153313	B2	20020521
US 2002-153831	A2	20020521
US 2002-153839	A2	20020521
US 2002-155373	A2	20020522
US 2002-155621	A2	20020522
US 2002-155703	A2	20020522
US 2002-155705	A2	20020522
US 2002-154594	A2	20020523
US 2002-154765	A2	20020523
US 2002-155097	A2	20020523
US 2002-412068P	P	20020918
US 2002-280315	A2	20021025
US 2002-302010	A2	20021121
US 2002-302614	A2	20021121
US 2002-322227	A2	20021217
US 2003-633876	A2	20030804
US 2003-633877	A2	20030804
US 2001-294203P	P	20010524
US 2001-296225P	P	20010605
US 2001-317479P	P	20010905
US 2001-335049P	P	20011030
US 2001-336218P	P	20011030
US 2001-345145P	P	20011109
WO 2002-US37491	W	20021121
US 2003-734902	A1	20031212
US 2003-735198	A1	20031212
US 2003-735199	A1	20031212
US 2003-735495	A1	20031212
US 2003-735496	A1	20031212
US 2003-735497	A1	20031212
US 2003-749535	A1	20031230
US 2003-749536	A1	20031230
US 2003-749537	A1	20031230
US 2003-749539	A1	20031230
US 2003-749783	A1	20031230
US 2003-750303	A1	20031230
US 2004-813721	A1	20040331
US 2004-816492	A1	20040401

AB The present invention provides novel condensation aerosols for the treatment of disease and/or intermittent or acute conditions. These condensation aerosols have little or no pyrolysis degradation products and are characterized by having an MMAD of between 1-3 μ . The aerosols are

made by rapidly heating a substrate coated with a thin film of drug having a thickness of between 0.05 and 20 μm , while passing a gas over the film, to form particles of a desirable particle size for inhalation. Kits comprising a drug and a device for producing a condensation aerosol are also provided. The device contained in the kit typically, has an element for heating the drug which is coated as a film on the substrate and contains a therapeutically ED of a drug when the drug is administered in aerosol form, and an element allowing the vapor to cool to form an aerosol. Also disclosed, are methods for using these aerosols and kits. For example, acebutolol (MW 336, m.p. 123°, oral dose 400 mg), a β -adrenergic blocker (cardiovascular agent), was coated on a stainless steel cylinder (8 cm). The drug (0.89 mg) was applied to the substrate, for a calculated drug film thickness of 1.1 μm . The substrate was heated at 20.5 V and purity of the drug aerosol particles was determined to be 98.9%; 0.53 mg was recovered from the filter after vaporization, for a percent yield of 59.6%. A total mass of 0.81 mg was recovered from the test apparatus and substrate, for a total recovery of 91%. High speed photographs were taken as the drug-coated substrate was heated to monitor visually formation of a thermal vapor. The photographs showed that a thermal vapor was initially visible 30 ms after heating was initiated, with the majority of the thermal vapor formed by 130 ms. Generation of the thermal vapor was complete by 500 ms.


IT 3605-01-4, Piribedil

RL: PEP (Physical, engineering or chemical process); PYP (Physical process); THU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses)


(drug condensation aerosols and kits for inhalation therapy)

RN 3605-01-4 CAPLUS

CN Pyrimidine, 2-[4-(1,3-benzodioxol-5-ylmethyl)-1-piperazinyl]- (9CI) (CA INDEX NAME)



L13 ANSWER 5 OF 19 CAPLUS COPYRIGHT 2006 ACS on STN
 AN 2004:131572 CAPLUS
 DN 140:292770
 TI Spectrophotometric Determination of Bisacodyl and Piribedil
 AU Abdel-Hay, Mohamed H.; Sabry, Suzy M.; Barary, Magda H.; Belal, Tarek S.
 CS Faculty of Pharmacy, Department of Pharmaceutical Analytical Chemistry,
 University of Alexandria, Alexandria, Egypt
 SO Analytical Letters (2004), 37(2), 247-262
 CODEN: ANALBP; ISSN: 0003-2719
 PB Marcel Dekker, Inc.
 DT Journal
 LA English
 AB Simple spectrophotometric methods are described for the assay of bisacodyl
 (BIS) and piribedil (PIR) based on charge-transfer and ion-pair
 complexation reactions. The 1st method is based on the reaction of the
 cited drugs with p-chloranilic acid (p-CA) in acetonitrile. The
 purple colored chromogen formed shows maximum absorbance at 518 nm.
 The 2nd method is concerned with the reaction of the investigated drugs
 with picric acid (PA) and 4 sulfonylphthalein acid dyes, namely; bromocresol
 green (BCG), bromocresol purple (BCP), bromophenol blue (BPB)
 and thymol blue (TB). The yellow ion-pair complexes formed show
 absorption spectra with maxima within the range from 400 to 415 nm. The
 stoichiometric ratio was found to be 1:1, for all complexation reactions
 examined, as calculated by the continuous variations method. Beer's law
 validation, accuracy, precision, limits of detection, limits of
 quantification, and other aspects of anal. merit are presented in the
 text. The proposed methods were applied for the determination of the analytes
 in their pure forms and in pharmaceutical preps. The results were
 in good agreement with those obtained by the official and reported
 methods.
 IT 3605-01-4, Piribedil
 RL: ANT (Analyte); ANST (Analytical study)
 (spectrophotometric determination of bisacodyl and piribedil)
 RN 3605-01-4 CAPLUS
 CN Pyrimidine, 2-[4-(1,3-benzodioxol-5-ylmethyl)-1-piperazinyl]- (9CI) (CA
 INDEX NAME)



RE.CNT 21 THERE ARE 21 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 6 OF 19 CAPLUS COPYRIGHT 2006 ACS on STN

AN 2003:678514 CAPLUS

DN 139:191440

TI Methods of treating or preventing a cardiovascular condition using a cyclooxygenase-1 inhibitor

IN Krul, Elaine S.

PA USA

SO U.S. Pat. Appl. Publ., 32 pp.

CODEN: USXXCO

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 2003162824	A1	20030828	US 2002-292255	20021112
PRAI	US 2001-331346P	P	20011112		
	US 2001-338291P	P	20011113		

OS MARPAT 139:191440


AB Methods for treating or preventing one or more cardiovascular conditions in a subject comprises treating the subject with a therapeutically effective amount of a selective cyclooxygenase-1 inhibitor or a pharmaceutically-acceptable salt, tautomer or prodrug thereof alone or in combination with either a drug used in the treatment or prevention of a cardiovascular condition or a non-drug therapy used in the treatment of a cardiovascular condition. Cyclooxygenase-1 inhibitor, 5-(4-Chlorophenyl)-1-(4-methoxyphenyl)-3-(trifluoromethyl)pyrazole (I), was prepared from 4'-chloroacetophenone and (4-methoxyphenyl)hydrazine hydrochloride. I inhibited development of atherosclerosis in cholesterol-fed apoE knockout mice.

IT 3605-01-4, Piribedil


RL: BSU (Biological study, unclassified); PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(peripheral vasodilator; cyclooxygenase-1 inhibitor for treating or preventing cardiovascular conditions)

RN 3605-01-4 CAPLUS

CN Pyrimidine, 2-[4-(1,3-benzodioxol-5-ylmethyl)-1-piperazinyl]- (9CI) (CA INDEX NAME)



L13 ANSWER 7 OF 19 CAPLUS COPYRIGHT 2006 ACS on STN
 AN 2003:304310 CAPLUS
 DN 139:47022
 TI Piribedil and bromocriptine in Parkinson's disease: A single-blind crossover study
 AU Tan, E. K.; Ratnagopal, P.; Han, S. Y.; Wong, M. C.
 CS Department of Neurology, Singapore General Hospital, Singapore, 169608, Singapore
 SO Acta Neurologica Scandinavica (2003), 107(3), 202-206
 CODEN: ANRSAS; ISSN: 0001-6314
 PB Blackwell Munksgaard
 DT Journal
 LA English
 AB Introduction: Clinicians switch from one dopamine agonist to another for various reasons. However, each change may inadvertently result in certain potential risks such as decreased medication efficacy or new side-effects. We evaluated the tolerability of a switch of bromocriptine to piribedil using two conversion ratios as a primary outcome measure, with motor function as a secondary outcome measure, in patients with mild to moderate Parkinson's disease (PD). Methods: Twenty consecutive patients with mild to moderate PD (Hoehn and Yahr, stage II-III) on treatment with stable doses of bromocriptine and levodopa were randomized to two groups of 10 patients each, to receive piribedil based on 1:5 or 1:10 conversion ratios. Blinded evaluations were performed: (1) United Parkinson's Diseased Rating Scale (UPDRS) scores both in "on" and "off", (2) Open-ended interviews for adverse events, (3) Epworth Sleepiness Scale, (4) Purdue Pegboard assessment during "on" and "off", (5) Hand-arm movement test during "on" and "off", and (6) Walking test during "on" and "off". Results: Major adverse events included "sleep attacks" in one patient and minor side-effects included giddiness, nausea, hallucinations, sleepiness and lethargy. However, these were mild and 19 (95%) of the 20 patients completed the study. There was a significant improvement in both the UPDRS "off" total and motor scores at 1 mo compared with baseline for the group on 1:10 ratio. The walking times during the "off" state at 1 and 2 mo were significantly better compared with baseline in the 1:5 group. There were otherwise no significant differences in the rating tests during both "off" and "on" states before and after the bromocriptine switch. Conclusions: We demonstrated that patients with mild to moderate PD who were on relatively low doses of bromocriptine can be safely switched to piribedil based on a conversion ratio of either 1:5 or 1:10. However, the higher conversion ratio has to be carried out with caution in patients with daytime somnolence.
 IT 3605-01-4, Trivastal
 RL: ADV (Adverse effect, including toxicity); PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses) (piribedil and bromocriptine treatment of Parkinson's disease)
 RN 3605-01-4 CAPLUS
 CN Pyrimidine, 2-[4-(1,3-benzodioxol-5-ylmethyl)-1-piperazinyl]- (9CI) (CA INDEX NAME)



L13 ANSWER 8 OF 19 CAPLUS COPYRIGHT 2006 ACS on STN

AN 2003:164583 CAPLUS

DN 139:127327

TI Electroanalytical characteristics of piribedil and its differential pulse and square wave voltammetric determination in pharmaceuticals and human serum

AU Uslu, Bengi; Ozkan, Sibel A.

CS Faculty of Pharmacy, Department of Analytical Chemistry, Ankara University, Ankara, 06100, Turk.

SO Journal of Pharmaceutical and Biomedical Analysis (2003), 31(3), 481-489
CODEN: JPBADA; ISSN: 0731-7085

PB Elsevier Science B.V.

DT Journal

LA English

AB The electrochem. oxidative behavior of piribedil (PR) was described. It was investigated by cyclic, linear sweep, differential pulse (DPV) and square wave (SWV) voltammetric techniques. The redox behavior of PR was found irreversible. Different parameters were tested to optimize the conditions for the determination of PR. The dependence of intensities of currents

and potential on pH, concentration, scan rate, nature of the buffer was investigated. Two sensitive methods for the measurement of PR were described. For anal. purposes, a very well resolved diffusion controlled voltammetric peak was obtained in 0.1 M H₂SO₄ and pH 5.7 acetate buffer. The determination peaks are obtained at 1.27 and 0.95 V for differential pulse and 1.29 and 0.97 V for SWV in 0.1 M H₂SO₄ and pH 5.7 acetate buffer, resp. The linear response was obtained in the ranges of 2×10^{-6} – 1×10^{-3} M in 0.1 M H₂SO₄ and 2×10^{-6} – 8×10^{-4} M in pH 5.7 acetate buffer for both techniques. The proposed techniques were successfully applied to the determination of PR in tablet dosage forms and human serum. Excipients did not interfere in the determination. The necessary statistical validation reveals that the proposed methods are free from significant systematic errors.


IT 3605-01-4, Piribedil

RL: ANT (Analyte); ANST (Analytical study)

(electroanal. characteristics of piribedil and its differential pulse and square wave voltammetric determination in pharmaceuticals and human serum)


RN 3605-01-4 CAPLUS

CN Pyrimidine, 2-[4-(1,3-benzodioxol-5-ylmethyl)-1-piperazinyl]- (9CI) (CA INDEX NAME)




RE.CNT 20 THERE ARE 20 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 9 OF 19 CAPLUS COPYRIGHT 2006 ACS on STN
 AN 2003:84772 CAPLUS
 DN 138:411186
 TI Piribedil-selective electrodes based on Bi(III)-iodide and Hg(II)-iodide complexes
 AU Abdel-Gawad, Fatama M.; Issa, Yousry M.; Hassouna, Mohamed E. M.; Hussien, Emad M.
 CS National Organization for Drug Control and Research, Giza, Egypt
 SO Mikrochimica Acta (2003), 141(1-2), 7-13
 CODEN: MIACAQ; ISSN: 0026-3672
 PB Springer-Verlag Wien
 DT Journal
 LA English
 AB The authors describe new ion-selective electrodes for piribedil (PD) based on Bi(III)-iodide or Hg(II)-iodide piribedil ion associate complexes in a poly(vinyl chloride) (PVC) membrane with dibutylphthalate (DBP) and dioctylphthalate (DOP) as plasticizing solvent mediator. The electrodes (PD-[BiI4]/DBP, PD-[BiI4]/DOP, PD-[HgI4]/DBP and PD-[HgI4]/DOP) show nearly Nernstian response over the concentration range 2×10^{-5} – 10^{-2} M of the drug. The working pH ranges were 3.1–6.0, 3.5–6.0, 3.0–6.0 and 3.5–6.4, and the isothermal coeffs. of the cells were 0.00113, 0.00146, 0.00059 and 0.00120 V/°C for PD-[BiI4]/DBP, PD-[BiI4]/DOP, PD-[HgI4]/DBP and PD-[HgI4]/DOP, resp. The selectivity coeffs. for numerous compds. are given. The electrodes were used for the direct determination of PD in pure form and in tablets or in biol. fluids by the standard addition and potentiometric titration methods.
 IT 3605-01-4, Piribedil
 RL: ANT (Analyte); ANST (Analytical study)
 (piribedil-selective electrodes based on Bi(III)-iodide and Hg(II)-iodide complexes for determination of piribedil in tablets and body fluids)
 RN 3605-01-4 CAPLUS
 CN Pyrimidine, 2-[4-(1,3-benzodioxol-5-ylmethyl)-1-piperazinyl]- (9CI) (CA INDEX NAME)




RE.CNT 14 THERE ARE 14 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 10 OF 19 CAPLUS COPYRIGHT 2006 ACS on STN
 AN 2001:263194 CAPLUS
 DN 135:111887
 TI Formulation and in vitro-in vivo evaluation of piribedil solid lipid micro- and nanoparticles
 AU Demirel, M.; Yazan, Y.; Muller, R. H.; Kilic, F.; Bozan, B.
 CS Faculty of Pharmacy, Department of Pharmaceutical Technology, Anadolu University, Eskisehir, 26470, Turk.
 SO Journal of Microencapsulation (2001), 18(3), 359-371
 CODEN: JOMIEF; ISSN: 0265-2048
 PB Taylor & Francis Ltd.
 DT Journal
 LA English
 AB Modification of the dissoln. rate and, thus, the enhancement of the bioavailability of a dopaminergic drug, piribedil, which has a low aqueous solubility and short elimination half-life have been the aim in this study. Prepsns. of micron and submicron particles using solid lipid carriers were performed for this purpose. For the avoidance of solvent residues resulting from the preparation technique, cold and hot homogenization methods were used to prepare solid lipid particles. After obtaining an appropriate particle size, piribedil loading and preparation yield by the use of those two methods, various formulations were prepared with different lipid, drug and surfactant materials. The factors mentioned affected properties of the particles, and the release rate was the fastest in acidic medium. Suspensions of pure piribedil and a formulation, selected according to the results obtained from in vitro dissoln. and particle size expts., were compared by using tremor tests in mice. The same suspensions were applied orally to rabbits and bioavailability of the solid lipid particle was found to be higher than the pure piribedil. After an in vitro-in vivo evaluation of piribedil solid lipid particles developed for Parkinson's disease therapy, it was determined that release rate was controlled and piribedil bioavailability could be improved.
 IT 3605-01-4, Piribedil
 RL: BPR (Biological process); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses) (formulation and in vitro-in vivo evaluation of piribedil solid lipid micro- and nanoparticles)
 RN 3605-01-4 CAPLUS
 CN Pyrimidine, 2-[4-(1,3-benzodioxol-5-ylmethyl)-1-piperaziny]- (9CI) (CA INDEX NAME)




RE.CNT 37 THERE ARE 37 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 11 OF 19 CAPLUS COPYRIGHT 2006 ACS on STN
 AN 2000:554279 CAPLUS
 DN 133:227937
 TI Poly(vinyl chloride) ion-selective electrodes for Piribedil determination
 AU Issa, Y. M.; Hassouna, M. M.; Abdel-Gawad, F. M.; Hussien, E. M.
 CS Faculty of Science, Cairo University, Giza, Egypt
 SO Journal of Pharmaceutical and Biomedical Analysis (2000), 23(2-3), 493-502
 CODEN: JPBADA; ISSN: 0731-7085
 PB Elsevier Science B.V.
 DT Journal
 LA English
 AB Piribedil (PD) ion-selective electrodes were constructed from poly(vinyl chloride) matrix membrane containing piribedil-tetraphenylborate (PD-TPB) as the electroactive component with dibutylphthalate or dioctylphthalate as the plasticizing solvent mediator. The electrodes displayed a linear response over the concentration range 2.0×10^{-5} to 10^{-2} M PD. The working pH ranges of the electrodes were 3.5-6.4 and 3.0-6.0, and the isothermal coeffs. of the cells were 0.00129 and 0.00096 V/°C, resp. The electrodes were used for the determination of the diprotonated PD species, the most successful being that based on dioctylphthalate solvent mediator. The electrodes show a linear response over the concentration range of 8.0×10^{-6} to 10^{-2} M PD, with Nernstian slope 30 mV/PD concentration decade when preconditioned by soaking in distilled water for 30 min. The electrodes exhibit good selectivity for the PD with respect to a large number of inorg. cations and organic substances of biol. fluids. Piribedil was determined successfully in pure solns. and in tablets or in biol. fluids using the standard addns. and potentiometric titration methods. The membrane withstood soaking in distilled water for more than 5 mo.
 IT 3605-01-4, Piribedil
 RL: ANT (Analyte); ANST (Analytical study)
 (determination of piribedil in tablets and biol. fluids by potentiometric titration using poly(vinyl chloride) ion-selective electrode)
 RN 3605-01-4 CAPLUS
 CN Pyrimidine, 2-[4-(1,3-benzodioxol-5-ylmethyl)-1-piperazinyl]- (9CI) (CA INDEX NAME)



RE.CNT 13 THERE ARE 13 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 12 OF 19 CAPLUS COPYRIGHT 2006 ACS on STN
 AN 1996:110574 CAPLUS
 DN 124:194199
 TI Effects of a dopaminergic agonist in the guinea pig cochlea
 AU d'Aldin, Christine; Puel, Jean-Luc; Leducq, Regine; Crambes, Olivier;
 Eybalin, Michel; Pujol, Remy
 CS Laboratoire de Neurobiologie de l'Audition et Plasticite Synaptique, CHU
 Hopital St. Charles, Montpellier, 34295, Fr.
 SO Hearing Research (1995), 90(1/2), 202-11
 CODEN: HERED3; ISSN: 0378-5955
 PB Elsevier
 DT Journal
 LA English
 AB This study investigated the role of dopamine, a putative lateral efferent
 neurotransmitter/modulator, in cochlear physiol. and physiopathol.
 Cochlear potentials were recorded in guinea pigs after intracochlear
 perfusion of increasing concns. (0.1-1 mM) of piribedil, an agonist of the
 D2/D3 receptors. A concentration-dependent reduction of the amplitude of
 auditory
 nerve compound action potential (CAP) was observed, predominantly at
 high-intensity tone-burst stimulations, and without significant effect on
 CAP threshold. There was no variation of cochlear microphonic and
 summing potential. When 1 mM piribedil was perfused into the cochlea
 during continuous 130-dB pure tone exposure (6 kHz, 15 min), CAP
 threshold shifts were less than in control animals with artificial
 perilymph-perfused cochleas. No dendritic damage was observed, although
 there was evident hair cell damage. Similarly, radial dendrites were
 clearly protected against ischemia-induced damage when 1 mM piribedil was
 applied prior to a 10-min ischemia. These results suggest that dopamine
 modulates the activity of radial afferent fibers via D2/D3 receptors. The
 protective effect of piribedil during acoustic trauma or ischemia suggests
 that this modulation corresponds to a prevention of excitotoxicity due to
 dysfunction of inner hair cell neurotransmission.
 IT 3605-01-4, Piribedil
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological
 study, unclassified); BIOL (Biological study)
 (cochlea response to)
 RN 3605-01-4 CAPLUS
 CN Pyrimidine, 2-[4-(1,3-benzodioxol-5-ylmethyl)-1-piperazinyl]- (9CI) (CA
 INDEX NAME)



L13 ANSWER 13 OF 19 CAPLUS COPYRIGHT 2006 ACS on STN

AN 1995:997477 CAPLUS

DN 124:146205

TI Method of manufacturing 2-[4-(1,3-benzodioxol-5-ylmethyl)-1-piperazinyl]pyrimidine

IN Chilmonczyk, Zdzislaw; Zaworska, Alicja; Cybulski, Jacek; Szelejewski, Wieslaw; Krajewski, Krzysztof; Dzikowska, Jadwiga

PA Instytut Farmaceutyczny, Pol.

SO Pol., 3 pp.

CODEN: POXXA7

DT Patent

LA Polish

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	PL 167397	B1	19950831	PL 1992-295856	19920908
PRAI	PL 1992-295856		19920908		

OS CASREACT 124:146205


AB Title compound I, also known as piribedil and useful as dopamine receptor agonist (no data), was prepared by reaction of 1-(2-pyrimidinyl)piperazine with piperonal in the presence of formic acid as reducing agent in 100-140°. This process results in 98% purity.

IT 3605-01-4P


RL: IMF (Industrial manufacture); PREP (Preparation)
(method of manufacturing 2-[4-(1,3-benzodioxol-5-ylmethyl)-1-piperazinyl]pyrimidine)

RN 3605-01-4 CAPLUS

CN Pyrimidine, 2-[4-(1,3-benzodioxol-5-ylmethyl)-1-piperazinyl]- (9CI) (CA INDEX NAME)



L13 ANSWER 14 OF 19 CAPLUS COPYRIGHT 2006 ACS on STN
 AN 1995:988974 CAPLUS
 DN 124:66420
 TI Effects of β -cyclodextrins on skin: implications for the transdermal delivery of piribedil and a novel cognition enhancing-drug, S-9977
 AU Legendre, J. Y.; Rault, I.; Petit, A.; Luijten, W.; Demuynck, I.; Horvath, S.; Ginot, Y. M.; Cuine, A.
 CS Ardix, Rue Eugene Vignat, Orleans, 45000, Fr.
 SO European Journal of Pharmaceutical Sciences (1995), 3(6), 311-22
 CODEN: EPSCED; ISSN: 0928-0987
 PB Elsevier
 DT Journal
 LA English
 AB The effects of β -cyclodextrin (β -CD), randomly methylated β -cyclodextrin (RAMEB) and 2-hydroxypropyl β -cyclodextrin (HP β -CD) on skin were investigated. The three cyclodextrins (CDs) were able to destabilize model liposomes and to extract significant amts. of cholesterol from isolated stratum corneum (SC). However, only RAMEB extracted all the major lipid classes from isolated SC, as shown by thin layer chromatog. Both RAMEB and HP β -CD could release 5-10% of the extractable cholesterol as well as proteins from hairless rat skin. Nevertheless, CDs did not induce any major modification of the differential scanning calorimetry (DSC) profile or the Fourier-transformed IR (FTIR) spectrum of SC. This was explained by the low percutaneous penetration of CDs. Furthermore, the influence of RAMEB on the transdermal diffusion through hairless rat skin of piribedil, a central dopaminergic agonist and of S-9977, a novel cognition enhancing drug, was studied. RAMEB was found to decrease the transdermal flux of piribedil, with which it forms an inclusion complex, as shown by NMR. Conversely, RAMEB increased by 2-fold the percutaneous absorption of the S-9977 hydrochloride, which does not interact with CD. Finally, a combination of oleic acid and RAMEB greatly increased by about 30-fold the flux of S-9977 hydrochloride.
 IT 3605-01-4, Piribedil
 RL: BPR (Biological process); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); PROC (Process); USES (Uses) (effects of β -cyclodextrins on skin: implications for the transdermal delivery of piribedil and a novel cognition enhancing-drug, S-9977)
 RN 3605-01-4 CAPLUS
 CN Pyrimidine, 2-[4-(1,3-benzodioxol-5-ylmethyl)-1-piperazinyl]- (9CI) (CA INDEX NAME)



L13 ANSWER 15 OF 19 CAPLUS COPYRIGHT 2006 ACS on STN

AN 1988:604820 CAPLUS

DN 109:204820

TI Reduced amounts of S-adenosylmethionine decarboxylase in the adrenal glands of rats following administration of piribedil or 2-deoxyglucose

AU Ekker, Marc; Sourkes, Theodore L.; Gabor, Ron

CS Fac. Med., McGill Univ., Montreal, QC, H3A 1A1, Can.

SO Biochemical Pharmacology (1988), 37(19), 3613-18

CODEN: BCPCA6; ISSN: 0006-2952

DT Journal

LA English

AB The activity of S-adenosylmethionine decarboxylase (SAM-DC) decreases in the adrenal gland of the rat following phys. stress, metabolic stress, or administration of dopamine agonists. Immunotitration studies with a serum against purified rat liver SAM-DC showed that the reduction in activity of the enzyme following administration of 2-deoxyglucose or piribedil was paralleled by a decrease in the amount of immunoreactive protein. There was no difference in the half-life of SAM-DC activity between piribedil-treated rats and controls. The properties of an extensively purified preparation of the adrenal enzyme resembled those of SAM-DC obtained from rat liver. It is suggested that the reduction in adrenal SAM-DC activity and protein content caused by stress is due to a reduction in the rate of synthesis of the enzyme.


IT 3605-01-4, Piribedil

RL: BIOL (Biological study)


(adrenal adenosylmethionine decarboxylase inhibition by)

RN 3605-01-4 CAPLUS


CN Pyrimidine, 2-[4-(1,3-benzodioxol-5-ylmethyl)-1-piperazinyl]- (9CI) (CA INDEX NAME)



L13 ANSWER 16 OF 19 CAPLUS COPYRIGHT 2006 ACS on STN
AN 1985:482187 CAPLUS
DN 103:82187
TI Neuro-active drugs in the regulatory system of sexual behavior of the male rat
AU Soulairac, A.; Soulairac, M. L.
CS Psychophysiol. Lab., Sainte-Anne Hosp., Paris, Fr.
SO Current Clinical Practice Series (1984), 26(Endorphins, Neuroregul. Behav. Hum. Reprod.), 179-200
CODEN: CCPSEZ; ISSN: 0168-6917
DT Journal
LA English
AB Use of neurotransmitter agonists and antagonists indicated that sex activity of male adult rats is regulated by catecholaminergic (dopaminergic and adrenergic) receptors. Ablation of neocortical areas in the brain resulted in marked disturbances in sex behavior. In rats bearing small or large cortical lesions, the alterations in sex behavior were completely reversed by caffeine [58-08-2]; partially reversed by amphetamine [300-62-9], L-dopa [59-92-7], and amineptine [57574-09-1]; but unchanged by testosterone [58-22-0] or nicotine [54-11-5]. Evidently, neural mechanisms are involved in sex activity. Results are discussed in relation to the pure physiol. elements of sex activity and libido.
IT 3605-01-4
RL: BIOL (Biological study)
(sex activity response to, receptor mechanism for)
RN 3605-01-4 CAPLUS
CN Pyrimidine, 2-[4-(1,3-benzodioxol-5-ylmethyl)-1-piperazinyl]- (9CI) (CA INDEX NAME)



L13 ANSWER 17 OF 19 CAPLUS COPYRIGHT 2006 ACS on STN
AN 1983:516654 CAPLUS
DN 99:116654
TI Sulpiride binding in rat striatum - effect of dopamine agonists and
sulfhydryl group reagents
AU Woodruff, G. N.; Freedman, S. B.
CS Dep. Physiol. Pharmacol., Univ. Southampton, Southampton, S09 3TU, UK
SO Acta Pharmaceutica Suecica (1983), (Suppl. 1, Dopamine Recept. Agonists
1), 118-29
CODEN: APSXAS; ISSN: 0001-6675
DT Journal
LA English
AB S-(-)-Sulpiride [23672-07-3] was bound to purified rat striatal
membranes with a dissociation constant of 7.4 nM and a maximum specific binding
capacity of 240 fmol/mg protein. Seventeen neuroleptics were compared for
their ability to displace the sulpiride binding. Generally, the potency
in displacing sulpiride binding was correlated with their clin.
effectiveness. The effect of 24 dopamine agonists for their ability to
displace sulpiride was also compared. Dopamine agonist affinity for the
sulpiride binding sites was generally decreased by the guanine nucleotide
Gpp(NH)p [34273-04-6]. Sulpiride binding was also inhibited by the
sulfhydryl reagent N-ethylmaleimide [128-53-0], indicating the presence
of an essential sulfhydryl group at the active site of the dopamine
receptor.
IT 3605-01-4
RL: BIOL (Biological study)
(sulpiride binding by brain striatum inhibition by)
RN 3605-01-4 CAPLUS
CN Pyrimidine, 2-[4-(1,3-benzodioxol-5-ylmethyl)-1-piperazinyl]- (9CI) (CA
INDEX NAME)



L13 ANSWER 18 OF 19 CAPLUS COPYRIGHT 2006 ACS on STN

AN 1983:448302 CAPLUS

DN 99:48302

TI Biphasic effects of some dopamine agonists on striatal acetylcholine concentrations

AU Waldmeier, Peter C.

CS Res. Dep., Ciba-Geigy Ltd., Basel, CH-4002, Switz.

SO European Journal of Pharmacology (1983), 90(1), 115-20

CODEN: EJPHAZ; ISSN: 0014-2999

DT Journal

LA English

AB Since the reduction of striatal dopaminergic transmission decreases striatal acetylcholine (ACh) [51-84-3] levels due to disinhibition of the resp. neurons, such an effect might be expected after selective stimulation of dopamine (DA) [51-61-6] autoreceptors. The effects of a number of DA agonists, including the purportedly selective presynaptic agents N-(propyl-3-(3-hydroxyphenyl)piperidine (3-PPP) [75240-91-4] and TL 99 [66543-77-9], on striatal ACh levels were investigated over a wide dose range in rats. Apomorphine [58-00-4] and 3-PPP decreased ACh levels in a lower dose range (0.01-0.03 mg/kg s.c. and 0.2-1 mg/kg s.c., resp.). TL 99 showed a much smaller effect (0.1-0.3 mg/kg s.c.), whereas piribedil [3605-01-4] and bromocriptine [25614-03-3] only increased ACh. However, 3-PPP (at 3 mg/kg and above) and TL 99 (at 3 mg/kg) increased ACh in much the same way as did conventional DA agonists. Apparently, pre- and postsynaptic DA receptors are distinct in a functionally relevant manner, and 3-PPP and TL 99 possess postsynaptic effects on DA receptors associated with cholinergic neurons. Since 3-PPP does not elicit stereotypies in spite of evidence for an involvement of cholinergic neurons in the mediation of this behavior it might be assumed that it acts on other postsynaptic DA receptors than does apomorphine. It seems possible that the 2 types of DA receptors are located on 2 different types of cholinergic neurons with different functions.


IT 3605-01-4

RL: BIOL (Biological study)


(acetylcholine of brain striatum response to)

RN 3605-01-4 CAPLUS

CN Pyrimidine, 2-[4-(1,3-benzodioxol-5-ylmethyl)-1-piperazinyl]- (9CI) (CA INDEX NAME)



L13 ANSWER 19 OF 19 CAPLUS COPYRIGHT 2006 ACS on STN
 AN 1981:473556 CAPLUS
 DN 95:73556
 TI Measurement of local partial oxygen pressure, thermal conductivity, and temperature of skeletal muscle in anesthetized dogs. Study on effects of different vasodilators
 AU Pourrias, Bernard; Tisne-Versailles, Jacky; Jammes, Michel
 CS Unite Pharmacol. Cardiovasc., Cent. Rech. Delalande, Rueil Malmaison, F 92500, Fr.
 SO Journal de Pharmacologie (1981), 12(2), 189-205
 CODEN: JNPBAG; ISSN: 0021-793X
 DT Journal
 LA French
 AB In the anesthetized dog, occlusion of the commons femoral artery caused a fall in pO₂, temperature, and thermal conductivity (index of muscular blood flow) in the stimulated or resting gastrocnemius muscle. In the stimulated muscle, the fall in pO₂ was very marked and the duration of hypoxia was longer than in the resting muscle. When the occlusion was released, the femoral flow increased, and the reactive hyperemia was greater when the muscle was stimulated. The variation of thermal conductivity, the kinetics of the decrease of pO₂ during arterial occlusion, and the amplitude of the reactive hyperemia may be an index of the development of collateral circulation. Piribedil [3605-01-4] (0.1 mg/kg, i.v.) increased femoral artery blood and i.m. thermal conductivity, temperature, and pO₂. Papaverine [58-74-2] (3 mg/g, i.v.) produced femoral vasodilation with inconsistent changes in i.m. thermal conductivity, temperature and pO₂. Intraduodenal administration of cinepazide maleate (I) [26328-04-1] (30 mg/kg) increased the arteriovenous O and glucose differences and decreased the fall in pO₂ and temperature during arterial occlusion. In also decreased the magnitude of the hyperemic reaction. Cinepazied acts at the local muscular level by interacting with adenosine and(or) by stimulating presynaptic purinergic receptors. Determination of the variations in local oxygen pressure, temperature, and thermal conductivity during an arterial occlusion and the hyperemic phase of the postocclusive reaction offers new possibilities for the screening of drugs for treating peripheral vascular disease.
 IT 3605-01-4
 RL: BIOL (Biological study)
 (muscle microcirculation response to, in artery occlusion)
 RN 3605-01-4 CAPLUS
 CN Pyrimidine, 2-[4-(1,3-benzodioxol-5-ylmethyl)-1-piperazinyl]- (9CI) (CA INDEX NAME)




```
=> s ethanol
L14      253287 ETHANOL

=> s l8 and l14
L15      6 L8 AND L14

=> s l15 not (l12 or l10)
L16      5 L15 NOT (L12 OR L10)

=> d l16 1-5 bib,ab,hitstr
```

LI6 ANSWER 1 OF 5 CAPLUS COPYRIGHT 2006 ACS on STN

AN 2004:59549 CAPLUS

DN 140:117387

TI Transdermal delivery of antiparkinson agents with skin penetration enhancer and volatile liquid

IN Klose, Kathryn Traci-Jane; Tran, Ngan Thi Kim; Morgon, Timothy Matthias; Finnin, Barrie Charles; Reed, Barry Leonard

PA Monash University, Australia; Acrux Dds Pty Ltd.

SO U.S. Pat. Appl. Publ., 8 pp., Cont.-in-part of U.S. Ser. No. 910,780.

CODEN: USXXCO


DT Patent

LA English

FAN.CNT 6

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 2004013620	A1	20040122	US 2003-428016	20030502
	US 6929801	B2	20050816		
	WO 9729735	A1	19970821	WO 1997-AU91	19970219
	W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU				
	RW: KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG				
	EP 1674068	A1	20060628	EP 2005-22951	19970219
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI				
	US 6299900	B1	20011009	US 1998-125436	19981218
	AU 9952589	A1	19991202	AU 1999-52589	19991001
	US 2002028235	A1	20020307	US 2001-910780	20010724
	US 6818226	B2	20041116		
PRAI	AU 1996-8144	A	19960219		
	WO 1997-AU91	W	19970219		
	US 1998-125436	A3	19981218		
	US 2001-910780	A2	20010724		
	AU 1997-17134	A3	19970219		
	EP 1997-904304	A3	19970219		
OS	MARPAT 140:117387				
AB	The present invention provides a transdermal drug delivery system which comprises: a therapeutically effective amount of an antiParkinson agent; at least one dermal penetration enhancer, which is a safe skin-tolerant ester sunscreen ester; and at least one volatile liquid The invention also provides a method for administering at least one systemic acting antiParkinson agent to an animal which comprises applying an effective amount of the antiParkinson agent in the form of the drug delivery system of the present invention. The addition of the sunscreen ester dermal penetration enhancer, octyl salicylate, surprisingly caused a marked increase (>15-fold) in the transdermal delivery of ropinirole across the skin (p<0.05). A topical spray contains 5 % volume/volume ropinirole, 5 % volume/volume octyl salicylate, and aqueous ethanol.				
IT	3605-01-4, Piribedil RL: BSU (Biological study, unclassified); PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses) (antiParkinson agent; transdermal delivery of antiparkinson agents with skin penetration enhancer and volatile liquid)				
RN	3605-01-4 CAPLUS				
CN	Pyrimidine, 2-[4-(1,3-benzodioxol-5-ylmethyl)-1-piperazinyl]- (9CI) (CA				

INDEX NAME)



L16 ANSWER 2 OF 5 CAPLUS COPYRIGHT 2006 ACS on STN

AN 1999:763835 CAPLUS

DN 132:26843

TI Compounds, compositions and methods for treating erectile dysfunction

IN Shoemaker, James D.

PA Saint Louis University, USA

SO PCT Int. Appl., 38 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 2

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9960985	A2	19991202	WO 1999-US11589	19990526
	WO 9960985	A3	20000217		
	W:	AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
	RW:	GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
	US 6124461	A	20000926	US 1998-84849	19980526
	AU 9943141	A1	19991213	AU 1999-43141	19990526
PRAI	US 1998-84849	A	19980526		
	WO 1999-US11589	W	19990526		

AB Vasoactive compds. are described for the treatment of erectile dysfunction and impotence. The compds. are reaction products of an anionic or neg. charged vasoactive or erection-inducing component and a cationic or pos. charged vasoactive or erection-inducing component. These components are combined as acids and bases to form an organic salt or ionically bonded compound. The compds. have advantageous solubility characteristics and efficacy.

A compound of the invention is combined with a pharmaceutical vehicle to form a composition which preferably includes an emulsifier. A local anesthetic and/or androgenic steroids may also be included. Compns. of the invention may also include more than vasoactive organic salt compound. The composition can be

advantageously formulated and administered to allow self-adjusted dosing, while minimizing or preventing overdosing. Phentolamine alprostadilate and papaverine alprostadilate, both existing as compds., not mixts., were prepared and formulated into pharmaceutical compns.


IT 3605-01-4, Piribedil

RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)


(phentolamine alprostadilate and papaverine alprostadilate compns. for treatment of erectile dysfunction)

RN 3605-01-4 CAPLUS


CN Pyrimidine, 2-[4-(1,3-benzodioxol-5-ylmethyl)-1-piperazinyl]- (9CI) (CA INDEX NAME)



L16 ANSWER 3 OF 5 CAPLUS COPYRIGHT 2006 ACS on STN
AN 1977:594937 CAPLUS
DN 87:194937
TI Alterations in dopamine receptor sensitivity by chronic ethanol treatment
AU Hoffman, Paula L.; Tabakoff, Boris
CS Med. Cent., Univ. Illinois, Chicago, IL, USA
SO Nature (London, United Kingdom) (1977), 268(5620), 551-3
CODEN: NATUAS; ISSN: 0028-0836
DT Journal
LA English
AB Mice fed a liquid diet containing 7% (volume) EtOH [64-17-5] for 7 days were, on withdrawal, less sensitive to the hypothermic effects of either EtOH (3 mg/kg, i.p.) or piribedil [3605-01-4] (20-60 mg/kg, i.p.), but as sensitive to clonidine [4205-90-7] (0.5 mg/kg, i.p.) as controls. Stimulation of adenylate cyclase (I) [9012-42-4] by dopamine [51-61-6], which was unaffected by acute EtOH, was reduced after withdrawal of EtOH from EtOH-fed mice. In addition, the I response to dopamine in EtOH-fed, withdrawn mice failed to reach the same maximum value as controls. The effects of EtOH withdrawal on EtOH tolerance and dopamine-sensitive I activity decreased with increasing time of withdrawal, becoming absent by 7 days after withdrawal. Tolerance to EtOH after chronic EtOH treatment appears therefore to result from an altered sensitivity of dopamine receptors.
IT 3605-01-4
RL: BIOL (Biological study)
(hypothermia from, ethanol effect on, dopamine receptors in relation to)
RN 3605-01-4 CAPLUS
CN Pyrimidine, 2-[4-(1,3-benzodioxol-5-ylmethyl)-1-piperazinyl]- (9CI) (CA INDEX NAME)



L16 ANSWER 4 OF 5 CAPLUS .COPYRIGHT 2006 ACS on STN
AN 1974:486051 CAPLUS
DN 81:86051
TI Suppression by dopamine agonists of the ethanol-induced stimulation of locomotor activity and brain dopamine synthesis
AU Carlsson, Arvid; Engel, Jorgen; Strombom, Ulf; Svensson, Torgny H.; Waldeck, Bertil
CS Dep. Pharmacol., Univ. Goeteborg, Goeteborg, Swed.
SO Naunyn-Schmiedeberg's Archives of Pharmacology (1974), 283(2), 117-28
CODEN: NSAPCC; ISSN: 0028-1298
DT Journal
LA English
AB The dopamine [51-61-6] receptor-stimulating stimulating agents apomorphine-HCl (I) [314-19-2] and ET 495 [3605-01-4], which had no marked effect on the locomotor activity, both inhibited the locomotor stimulation induced by 2.2 g ethanol [64-17-5]/kg i.p. in mice. I and ET 495 did not change the brain dopamine-3H and noradrenaline-3H contents of mice injected with tyrosine-3H. However, both of them inhibited the EtOH-induced increased in net yield of dopamine-3H. This effect of I and ET 495 could not be ascribed to changes in the specific activity of tyrosine-3H in the plasma, and I had no effect on the blood level of EtOH as measured 1 hr after the administration of EtOH. The possibility that these effects of the dopamine agonists may be mediated by stimulation of presynaptic, inhibitory receptors is discussed.
IT 3605-01-4
RL: BIOL (Biological study)
(ethanol-induced brain dopamine formation and locomotor activity in response to)
RN 3605-01-4 CAPLUS
CN Pyrimidine, 2-[4-(1,3-benzodioxol-5-ylmethyl)-1-piperazinyl]- (9CI) (CA INDEX NAME)



L16 ANSWER 5 OF 5 CAPLUS COPYRIGHT 2006 ACS on STN

AN 1965:480704 CAPLUS

DN 63:80704

OREF 63:14880e-h,14881a

TI Pyrimidine derivatives

PA Science Union et Cie., Societe Francaise de Recherche Medicale

SO 21 pp.

DT Patent

LA Unavailable

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	NL 6413349		19650520	NL 1964-13349	19641117
PRAI	GB		19631113		

AB I and their salts show vasodilation effects and are useful as analgetics and antiinflammatory agents. To 21 g. N-(3,4-methylenedioxybenzyl)piperazine (II) in 300 ml. dry xylene was added 28 g. K₂CO₃ and 11.3 g. 2-chloropyrimidine (III), the suspension refluxed 9 hrs., the mixture extracted with 10% HCl, the acid layer washed with Et₂O and made alkaline (K₂CO₃), the separated oil extracted with CHCl₃, the CHCl₃ dried (K₂CO₃)

and evaporated, and the oil (20 g.) crystallized from EtOH to give 15 g. I (R = R₁

= R₃ = H, R₂ = 3,4-methylenedioxybenzyl) (IV), m. 96°. Heating a mixture of 23.2 g. II, 12.5 g. III, 31 g. K₂CO₃, and 150 ml. HCONMe₂ 8.5 hrs. at 130°, the solvent evaporated in vacuo, and the hot residue poured into 100 ml. boiling H₂O gave 28 g. IV, m. 97-8° (EtOH).

The following I (R = R₁ = H) were similarly prepared (R₂, R₃, m.p., salt, and its m.p. given): 3,4-ethylenedioxybenzyl, H, -, di-HCl, 220-6° (decomposition); Ph₂CH, H, 170°, -, -, 2,3,4-(MeO)₃C₆H₂CH₂, H, 105°, -, -, 2-MeOC₆H₄, H, 73°, -, -, 3-MeOC₆H₄, H, 78°

-, -, 4-MeOC₆H₄, H, 108-10°, -, -, 3,4-(MeO)₂C₆H₃CH₂, 101°,

-, -, 2,3-(MeO)₂C₆H₃CH₂, -, di-HCl, 207-15° (decomposition);

3,4-methylenedioxybenzyl, 3-Me, -, di-HCl; 189-97° (decomposition);

3,4-methylenedioxybenzyl, 2-Me, -, di-HCl, 225-8° (decomposition);

3,4-MeO(OH)C₆H₃CH₂, H, -, di-HCl hydrate, 180-8°; 3,4(HO)₂C₆H₃CH₂,

H, -, di-HCl hydrate, 207-12° (decomposition). The following I (R₂ =

3,4-methylenedioxybenzyl, R₃ = H) were also prepared (R, R₁, m.p., salt, and

its m.p. given): 4-MeO, H, 89-90°, -, -, 4-EtO, H, -, HCl,

225°; 4-Me, 6-Me, -, HCl, 256°; 4-Me, 5-Me, -, HCl,

245°; 4-Me, H, -, di-HCl, 212-15° (decomposition); 4-NH₂, H,

169°, -, -, 4-NHMe, H, -, bis(methanesulfonate), 234°;

4-NMe₂, H, 103°, -, -, 4-N-(CH₂CH₂OH)₂, H, -, di-HCl,

216-25° (decomposition); 4-OH, H (V), 214°, -, -, 4-OCH₂Ph, H

(VI), 108°, -, -. Hydrogenation of VI (10% Pd-C, 15 atmospheric) gave V.


IT 3605-01-4, Pyrimidine, 2-(4-piperonyl-1-piperazinyl)-

(preparation of)

RN 3605-01-4 CAPLUS

CN Pyrimidine, 2-[4-(1,3-benzodioxol-5-ylmethyl)-1-piperazinyl]- (9CI) (CA

INDEX NAME)



10/810,799

=> log y

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

136.36

304.83

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

SINCE FILE

TOTAL

ENTRY

SESSION

CA SUBSCRIBER PRICE

-18.75

-18.75

STN INTERNATIONAL LOGOFF AT 15:31:44 ON 21 AUG 2006